

2. 14645-53477-2-CE.docx

anonymous marking enabled

Submission date: 16-Mar-2025 01:17PM (UTC-0700)

Submission ID: 2498610033

File name: 2_14645-53477-2-CE.docx (602.32K)

Word count: 4857

Character count: 28995



Antioxidant of Ethanol Extract and Toxicity of Fractions from *Aspergillus unguis* a Marine Sponge Symbiont of *Aaptos suberitoides*

Scify Bilqis Nawafi Masyerli¹, Mai Efdi¹, Muhammad Hasan Bashari², Mochamad Untung Kurnia Agung³, Beginer Subhan⁴, Efahmi^{5,6}, Yosie Andriani⁶, Syafrizayanti^{1*}

¹ Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Andalas, Padang, West Sumatera, Indonesia

² Department of Biomedical Sciences, Faculty of Medicine, Universitas Padjadjaran, Bandung, West Java, Indonesia

³ Department of Marine Science, Faculty of Fisheries and Marine Science, Universitas Padjadjaran, Bandung, West Java, Indonesia

⁴ Department of Marine Science and Technology, Faculty of Fisheries and Marine Science, IPB University, Bogor, West Java, Indonesia

⁵ School of Pharmacy, Bandung Institute of Technology, Bandung, West Java, Indonesia

⁶ University Center of Excellence for Nutraceuticals, Bioscience and Biotechnology Research Center, Bandung Institute of Technology, Bandung, West Java, Indonesia

* Corresponding Author e-mail: syafrizayanti@sci.unand.ac.id

Article History

Received: 03-01-2025

Revised: 24-02-2025

Published: 28-02-2025

Keywords:

antioxidant; *Aspergillus unguis*; brine shrimp lethality test; fractions; toxicity

Abstract

The marine sponge *Aaptos suberitoides* has been shown to have anticancer properties, with clear evidence of its capacity to suppress the growth of cancer cells. However, the pharmaceutical exploration of chemicals from marine organisms causes significant environmental concerns. *Aspergillus unguis*, a fungal symbiont of the marine sponge *A. suberitoides*, has been isolated for its potential in sustainable natural products resources. This study assesses the antioxidant activity of ethanol extract and the toxicity of four levels fractions from ethanol extracts of *A. unguis* mycelium and these results were reported for the first time in this study. The antioxidant of ethanol extract was determined using the 2,2-Diphenyl-1-picrylhydrazyl (DPPH) method. The fractions were examined for toxicity using the Brine Shrimp Lethality Test (BSLT). The ethanol extract of *A. unguis* shows very strong antioxidant bioactivity ($IC_{50} = 42.84$ mg/L). The LC_{50} values for hexane, chloroform, ethyl acetate, and butanol fractions were determined to be 74.11 μ g/mL, 93.84 μ g/mL, 59.37 μ g/mL, and 142.79 μ g/mL, respectively. It indicates significant toxicity. These preliminary results are important knowledge for further research into the bioactivity potential of the metabolites as candidate anticancer compounds, aligning with marine pharmaceutical drug development.

How to Cite: Masyerli, S., Efdi, M., Bashari, M., Untung Kurnia Agung, M., Subhan, B., Efahmi, E., Andriani, Y., & Syafrizayanti, S. (2025). Antioxidant of Ethanol Extract and Toxicity of Fractions from *Aspergillus unguis* a Marine Sponge Symbiont of *Aaptos suberitoides*. *Hydrogen: Jurnal Kependidikan Kimia*, 13(1), 12-21. doi:<https://doi.org/10.33394/hjkk.v13i1.14645>



<https://doi.org/10.33394/hjkk.v13i1.14645>

This is an open-access article under the CC-BY-SA License.



26

INTRODUCTION

Cancer is a significant challenge in society, public health, and the economy in the 21st century, accounting for nearly one in six deaths (16.8%) globally and one in four deaths (22.8%) from noncommunicable diseases (NCDs). It is responsible for 30.3% of premature deaths from NCDs among individuals aged 30–69 years and ranks among the top three causes of death in this age group across 177 out of 183 countries (Bray Bsc et al., 2024; Bayani et al., 2024). Based on data from the Institute for Health Metrics and Evaluation (IHME) in 2019, cancer is the

second most common cause of death in Indonesia, accounting for 13.4% of total deaths, following coronary heart disease (38.2%) and preceding diabetes (8.7%) and chronic obstructive pulmonary disease (COPD) (5.9%) (Andinata et al., 2023). Cancer continues to be a leading cause of mortality globally, requiring ongoing research into effective therapeutic agents (Min & Lee, 2022).

Natural sources, including marine organisms, offer a rich repository for the discovery of novel anticancer compounds due to their high biodiversity and the diversity of bioactive compounds (Santaniello et al., 2023). According to the latest statistics, marine-derived fungi contributed nearly half (47%) of all newly reported marine natural products (NPs) in 2019 (Van Anh et al., 2021). Notably, the marine sponge *Aaptos suberitoides* has been identified in multiple studies as a promising source of anticancer compounds (Yun et al., 2019).

A study has demonstrated that the ethanol extract of *A. suberitoides* exhibits cytotoxic properties, effectively inhibiting the proliferation and migration of the HCC-1954 cell line, a HER2+ breast cancer cell line resistant to trastuzumab—at a concentration of 12 µg/mL (Bashari et al., 2021). From the same species collected in Ambon, Indonesia, the alkaloid compounds, including aaptamine (8,9-dimethoxy-1H-benzo[de][1,6]naphthyridine), and four other derivatives that have been isolated and studied for anticancer properties (Pham et al., 2013). Several investigations have confirmed the bioactivity of aaptamine within the concentration range of 1–10 µM, showing cytotoxic effects on T47D breast cancer cells, murine cancer cells, and MG63 human osteosarcoma cells, by enhancing the regulation of p21 promoter expression and inducing cell cycle arrest (Aoki et al., 2006; Dyshlovoy et al., 2014; Tailor et al., 2024).

While the potential of marine organisms as sources for anticancer drug development is significant, the sustainability of marine ecosystems must be considered. Overexploitation of marine resources, including *A. suberitoides*, not only threatens marine biodiversity but also the sustainability of the organisms themselves (Varijakzhan et al., 2021). This dilemma highlights the necessity of marine conservation, which might be mitigated by exploring alternative sources such as microbial symbionts residing within marine organisms. These symbionts retain the capability to produce similarly significant bioactive compounds as their host organism (Turon et al., 2019). Among them, the genus *Aspergillus* is one of the most widespread filamentous fungi and serves as a key source of marine fungal natural products. This genus has been found to produce a wide array of structurally diverse secondary metabolites, including polyketides, alkaloids, terpenes, steroids, and peptides, many of which exhibit significant biological activities (Van Anh et al., 2021).

One example of a fungal species belonging to the genus *Aspergillus* is the marine sponge symbiont fungus *Aspergillus unguis*, associated with *A. suberitoides* collected from Harapan Island in Kepulauan Seribu, Indonesia (106°34'51.052" E, 5°38'55.462" S), shows considerable promise as a source of two novel anticancer compounds (Bashari et al., 2025). However, studies on the potential bioactivity investigation of extract and/or fractions of *A. unguis* as a source of compounds for anticancer potential candidates are still very limited and less explored. Therefore, the aim of this preliminary study is to investigate more potential from *A. unguis*, the antioxidant of ethanol extract and the toxicity of fractions from the ethanol extract of *A. unguis* for further emphasizing the value of symbiotic marine microbes in the development of potential candidate compounds for anticancer therapies.

METHOD

Materials

The experimental materials include isolate *A. unguis* from Mr. Mochamad Untung Kurnia (43), M.Si (the Marine Sciences Program, Universitas Padjadjaran), sterilized seawater, Potato Dextrose Agar (PDA), Potato Dextrose Broth (PDB), 70% alcohol (68), ethanol, hexane, chloroform, ethyl acetate, and butanol. Additional materials used are Whatman No. 42 filter paper, DPPH (2,2-diphenyl-1-picrylhydrazyl), ascorbic acid, 1% dimethyl sulfoxide (DMSO), and brine shrimp larvae (*Artemia salina* Leach).

Equipment and Instruments

The equipment utilized comprises general laboratory glassware, an inoculation loop, test tube racks, vial bottles, Pasteur pipettes, a laboratory shaker, and a magnetic stirrer. Precision instruments include an analytical balance (KERN ABS 220-4N), a laminar airflow cabinet (BioLab BLHZ-204), an incubator (Gallenkamp), a rotary evaporator (BUCHI R-100), and a UV-Vis spectrophotometer (Shimadzu UV-1280).

Procedures

Cultivation *A. unguis*

The isolate of the *A. unguis* was cultivated on Potato Dextrose Agar (PDA) for three days at room temperature (25-28°C). Then, culture was transferred to Potato Dextrose Broth (PDB) and maintained in 120 rpm for 30 days at 25-28°C. Each day, a (65) mL sample of the culture was extracted, diluted in a 1:1000 ratio with sea water, and its optical density measured at a wavelength of 405 nm using UV-VIS spectrophotometry, a procedure repeated twice following the methodology previously established (Nevalainen et al., 2014).

Extraction and Fractionation *A. unguis*

Approximately 5 liters of *A. unguis* culture were harvested on day 15 of incubation by filtering the mycelium using Whatman No. 42 filter paper, subsequently soaked in 500 mL of ethanol (repeated $\times 3$ or till exhaustion) over a three-day period with continuous shaking. Then the ethanol extract evaporated under a rotary evaporator and subsequently into freeze-drying till dryness. The fractionation of this dry extract began with hexane (repeated $\times 3$ or till exhaustion), followed in sequence by chloroform, ethyl acetate, and butanol. Each fraction of *A. unguis* was evaporated under a rotary evaporator. The dry fractions were stored at 4°C for subsequent experiments (Clark, James H., et al., 2017).

Antioxidant Assay of *A. unguis* Ethanol Extract Using DPPH Method

Two mL of ethanol extract solution with concentration were prepared in variations from 0 to 100 mg/L, and mixed with 3 mL of (63) DPPH solution (0.1 mM) and then incubated for 30 minutes in dark conditions. Ethanol served as the control, and ascorbic acid was used as a reference in (63) antioxidant testing. Subsequently, absorbance was measured using a UV-Vis spectrophotometer (Shimadzu UV-1280, Japan) at a wavelength of 517 nm, then plotted as a percentage of radical scavenging to calculate the IC₅₀ (Baliyan et al., 2022).

Toxicity Assay of *A. unguis* Fractions Using BSLT Method

Each fraction was tested for toxicity using the Brine Shrimp Lethality Test (BSLT). One gram of *Artemia salina* Leach eggs was hatched by soaking in 1 L of seawater with aeration in a glass box that was darkened on one side and illuminated with a 40-60 watt incandescent bulb for 48 hours. The hatched shrimp larvae migrated towards the brighter side of the glass box. Test solutions were prepared at various concentrations (0, 50, 100, 150, 250, 300 µg/mL) using the same solvent as the fraction. Control solutions were made without the addition of the

extract. 2 mL of each test solution was placed into a 30 mL vial bottle and dried until the solvent had completely evaporated. Then, 20 μ L of DMSO and 5 mL of seawater were added. Subsequently, 5 shrimp larvae were introduced into each bottle. Each concentration was tested in triplicate and compared with the control solution. Observations were made after 24 hours. The number of surviving shrimp larvae was counted and the LC_{50} value of each fraction was determined (Zakwan et al., 2023).

RESULTS AND DISCUSSION

Growth Curve of *A. unguis*

The growth of the fungus *A. unguis* is depicted in Figure 1, which illustrates four growth phases over a 21-day culture period: the lag or adaptation phase, the log or exponential phase, the stationary phase, and the death or decline phase. The lag phase occurs from day 1 to day 2, the log phase from day 3 to day 9, the stationary phase from day 10 to day 19, and the death phase begins on day 20 and continues thereafter. The harvest of *A. unguis* is timed at the 14-day mark of incubation during the stationary phase, where the production of secondary metabolites peaks (Oliveira et al., 2024). During the stationary phase, cell growth slows due to nutrient limitations, accumulation of metabolic waste products, and environmental stress, which stimulate the active and increased production of secondary metabolites as a form of adaptation and response mechanism (Molnár et al., 2010).




Figure 1. Growth curve of *A. unguis*

During the lag phase, fungi undergo adaptation to their environment. Fungal cells remain inactive in terms of division, while metabolic activity rises in preparation for subsequent stages. Essential molecules, including enzymes, are synthesized to facilitate growth. Following this adaptation, fungi enter the exponential or growth phase, where cellular division accelerates. At this stage, fungal growth peaks, and the cell population expands. The availability of nutrients and favorable environmental conditions supports this rapid growth (Stange et al., 2019).

However, once the optimal growth conditions are exhausted, fungi transition into the stationary phase, where growth stabilizes due to nutrient depletion and the buildup of metabolic waste. During this phase, fungal growth is balanced by the death of cells. It is also during this phase that fungi may begin to produce spores or other forms of resistance to cope with adverse conditions. Although carbon, a vital energy and nutrient source, is depleted, growth continues as dead cells undergo lysis and provide a recycled nutrient source (Oliveira et al., 2024). Ultimately, the death phase ensues, characterized by a significant reduction in biomass. In this

phase, the number of dead cells surpasses that of living cells, as nutrient availability becomes severely limited and waste product accumulation increases, resulting in a decline in fungal growth and eventual cell death (Rendowaty et al., 2017).

Moreover, growth conditions influence the growth phases; for example, the growth curve of *Aspergillus unguis* (WR8), symbiont of the sponge *Haliclona fascigera*, shows different phases when cultivated statically for 28 days and under shaking at 120 rpm for 14 days in Sabouraud Dextrose Broth (SDB). *Aspergillus unguis* (WR8) experiences its log phase from day 7 to day 14, the stationary phase from day 15 to day 21, and the death phase begins after day 21, making day 21 the optimal cultivation time for *A. unguis* (WR8). In contrast, under shaking conditions, *A. unguis* (WR8) only exhibits a log phase from day 2 to day 14 (Nursid et al., 2015; Rendowaty et al., 2017).

Extract and Fractions of *A. unguis*

The yield data for the extracts obtained from the maceration process using ethanol, as well as the weights of the *A. unguis* fractions, are presented in Table 1. The ethyl acetate fraction yielded the highest percentage (2.17%) compared to the other fractions. The extraction and fractionation processes were successfully conducted, resulting in yields that are considered substantial for fungal samples. Previous research has shown that the extraction of *A. unguis* (WR8), a symbiont of the sponge *Haliclona fascigera*, using a shaker resulted in a larger fungal biomass (1.45 g) due to stirring or agitation at 120 rpm, which rapidly increased fungal growth, while a smaller biomass (1.04 g) was observed under static conditions (Rendowaty et al., 2017).

Table 1. Amount and yield percentage of extract and fractions of *A. unguis*

| Extract/Fraction | Weight (gram) | Yield (%) |
|------------------|---------------|-----------|
| Ethanol | 72.5924 | 1.45 |
| Hexane | 1.2469 | 1.72 |
| Chloroform | 0.8752 | 1.21 |
| Ethyl Acetate | 1.5788 | 2.17 |
| Butanol | 1.3011 | 1.79 |

Agitation affects nutrient mixing, mass and heat transfer, changes in fungal morphology, variations in growth and metabolic product formation, and cell structure damage (Nursid et al., 2015; Rendowaty et al., 2017). Factors influencing the success of the extraction and fractionation processes include the methods and solvents used, environmental conditions such as temperature, sample size, media composition, culture conditions, and purification methods (Agarwal, 2018; Badar et al., 2022; Egbuna et al., 2020).

Antioxidant of *A. unguis* Etanol Extract

Nowadays, antioxidants have gained significant attention due to their beneficial effects as health promoters in the management of cardiovascular diseases, cancer, and aging, among other conditions. The growing global interest in identifying antioxidant compounds from natural sources has prompted us to evaluate the antioxidant activity of *A. unguis* extracts (Oogarah et al., 2020). The ethanol extract of *A. unguis* demonstrated the ability to scavenge PPH radicals, as shown in Figure 2.a, with ascorbic acid—a synthetic antioxidant—used as a reference in Figure 2.b. The scavenging activity of *A. unguis* ethanol extract against free radicals increased with higher concentrations, reaching an IC₅₀ of 42.84 mg/L, while ascorbic acid had an IC₅₀ of 5.18 mg/L. An extract is considered to possess very strong antioxidant bioactivity if its IC₅₀ value is less than 50 mg/L, and strong if the value is between 50 and 100 mg/L (Fristiody et al., 2020). Additionally, fungi that are symbiotic with marine sponges are found to exhibit bioactivity potential comparable to their sources, underscoring the significant therapeutic potential of marine-derived fungi.




Figure 2. (a) Antioxidant activity of *A. unguis* ethanol extract and (b) Standard antioxidant ascorbic acid

There is a limited amount of research on the antioxidant effects of sponges and/or its symbiont extracts. According to existing literature, several metabolites isolated from marine sponges, including indole derivatives, aromatic alkaloids, aromatic polyketides, and phenolic compounds, have demonstrated significant antioxidant activity, surpassing that of vitamin E and ascorbic acid. A scientific study identified alkaloids, phenols, steroids, terpenoids, tannins, and saponins in extracts from the sponge *Plakortia nigra* collected from the waters of Mauritius, with total phenolic content (TPC) ranging from 2.28 ± 0.072 to 12.79 ± 0.236 mg gallic acid equivalents per gram of extract. Similarly, the findings of this study align with previously reported occurrences of phenolic compounds in extracts of *Stylissa* sp. and *Biemna tubulosa* from Mauritius, with varying polarities (Oogarah et al., 2020).

The antioxidant bioactivity of the ethanol extract of *A. unguis* was first reported in this study. A previous study reported the methanol extract from *A. suberitoides*, the marine sponge originating as a symbiont of *A. unguis*, exhibited prominent antioxidant activity with an IC_{50} of $27.42 \mu\text{g/mL}$ (Abdillah et al., 2013). Similar studies on *Aaptos* sp., utilizing acetone as a solvent, demonstrated antioxidant properties with an IC_{50} of $16.10 \mu\text{g/mL}$ (Fristiohy et al., 2020). The ethyl acetate extract from *A. suberitoides* was found to be equivalent to $286 \mu\text{g}$ of ascorbic acid per $500 \mu\text{g/mL}$ of extract, indicating strong antioxidant potential (Rivera & Uy, 2012).

Additionally, *A. unguis* sourced from diverse symbionts such as the green microalga *Enteromorpha* sp. showed a moderate antioxidant activity at $73.91 \pm 0.91 \mu\text{g/mL}$, contrasting with the standard ascorbic acid which exhibited an activity of $28.01 \pm 0.5 \mu\text{g/mL}$ (Sajna et al., 2020). Another symbiotic variant, *Aspergillus unguis* associated with *Agelas* sp., inhibited acetylcholinesterase by 47.5% at a concentration of 400 g/mL and α -glucosidase by 42.3%, compared to the standard acarbose with 25% inhibition, showcasing its potential in enzymatic inhibition (Abd et al., 2015).

Toxicity of *A. unguis* Fractions

The ability of each fraction of *A. unguis* ethanol extract to induce mortality in *Artemia salina* Leach larvae is depicted in Figure 3. Determination of the toxicity of compounds or extracts acutely using shrimp larvae *Artemia salina* is a simple preliminary/prescriptive test of biological activity. This BSLT method can be used as a prescriptive test in the study of compounds that lead to cytotoxic activity tests. Each fraction exhibits varying levels of larvicidal activity, with the fastest responses and highest mortality rates occurring at a concentration of $300 \mu\text{g/mL}$. The higher the concentration of each fraction, the greater the

number of shrimp larvae that succumb. The LC_{50} values for each *A. unguis* fraction were determined by substituting the probit value of 50 into the linear regression equation $y = ax + b$, using Microsoft Excel. The calculated toxicity results for the *A. unguis* fractions reveal varying LC_{50} values, as illustrated in Figure 4.




Figure 3. Percentage of mortality of *A. unguis* fractions (a) Butanol (b) Ethyl acetate (c) Chloroform (d) Hexane

The toxicity values for the hexane, chloroform, ethyl acetate, and butanol fractions of *A. unguis* were determined to be 74.11 $\mu\text{g/mL}$, 93.84 $\mu\text{g/mL}$, 59.37 $\mu\text{g/mL}$, and 142.79 $\mu\text{g/mL}$, respectively. According to the toxicity classification criteria, a compound is considered highly toxic if its LC_{50} is less than 30 $\mu\text{g/mL}$, toxic if the LC_{50} is between 30 and 1000 $\mu\text{g/mL}$, and non-toxic if the LC_{50} exceeds 1000 $\mu\text{g/mL}$ (Zakwan et al., 2023). Based on these LC_{50} values, the fractions from *A. unguis* are categorized within the toxic range. These preliminary test results serve as a benchmark for further research into the bioactive potential of the metabolites contained in each fraction, particularly their anticancer properties. This study underscores the importance of evaluating the toxicological profiles of natural extracts in the development of therapeutic agents (Yun et al., 2019).

The toxicity levels of fractions derived from *A. unguis* have not been previously reported. However, the toxicity of the ethyl acetate extract from the sponge *A. suberitoides*, tested using the Brine Shrimp Lethality Test (BSLT), has been documented with an LC_{50} of 18.66 $\mu\text{g/mL}$. Fractionation of this ethyl extract through column chromatography revealed that the fourth ethyl acetate sub-fraction contains the bisdemethylaaptamine compound, an alkaloid secondary metabolite, with an LC_{50} of 2.56 $\mu\text{g/mL}$ (Situmorang, Fronika, 2012). The toxicity of the acetone extract of *Aaptos* sp. sponge was also reported, with an LC_{50} value of 1041.5 $\mu\text{g/mL}$, which is classified as non-toxic (Fristiody et al., 2020). The LC_{50} values are typically

obtained through statistical analysis, such as probit regression or log-logit analysis, based on the relationship between sample concentration and larval mortality percentage. A lower LC_{50} value indicates a higher toxicity of the tested sample (Fristiohady et al., 2020).




Figure 4. Fractions toxicity (LC_{50}) of *A. unguis*

This provides a basis for further research and evaluation of the potential toxicological impacts of these natural extracts, underscoring the critical role of dose-response assessments in the development of both therapeutic agents and toxicity studies.

CONCLUSION

Based on the search findings, it was discovered the ethanol extract of *A. unguis* has demonstrated very strong antioxidant activity with an IC_{50} value of 42.84 mg/L . Toxicity testing of the fractions from the ethanol extract of *A. unguis* indicates that all fractions are classified as toxic, with respective LC_{50} values of hexane (74.11 $\mu\text{g/mL}$), chloroform (93.84 $\mu\text{g/mL}$), ethyl acetate (59.37 $\mu\text{g/mL}$), and butanol (142.79 $\mu\text{g/mL}$). Therefore, the extract and its fractions from *A. unguis* show great potential for its toxicity and warrant further investigation for their anticancer capabilities against various cancer cells, which could lead to the exploration of their secondary metabolites for cancer drug development. All these investigation results related to the antioxidant of ethanol extract and the toxicity of four levels fractions from the ethanol extract of *A. unguis* marine symbiont sponge *A. suberitoides* are reported for the first time in this study.

RECOMMENDATIONS

The fractions of *A. unguis* can be further studied for their inhibitory abilities and anticancer mechanisms against various cancer cells. Additionally, the mechanisms of interaction between the compounds within these fractions and cancer target proteins can be examined to potentially develop them into structure-based drug discovery.

ACKNOWLEDGEMENTS

The authors express their gratitude to the Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Andalas, for providing the facilities for this study.

BIBLIOGRAPHY

- Abd, F. K., Hady, E., Shaker, K., & Elshahid, Z. A. (2015). Antioxidant, Acetylcholinesterase and α -glucosidase potentials of metabolites from the marine fungus *Aspergillus unguis* RSPG_204 associated with the sponge (*Agelas* sp.) GC-MS Analysis of Ethanol Extract of *Solanum* Species and *Ficus* Varieties from Saudi Arabia and their Systematics Implications View project. In *Article in International Journal of Pharmaceutical Sciences Review and Research* (Vol. 30, Issue 1). www.globalresearchonline.net
- Abdillah, S., Nurhayati, A. P. D., Nurhatika, S., Setiawan, E., & Heffen, W. L. (2013). Cytotoxic and antioxidant activities of marine sponge diversity at Pecaron Bay Pasir Putih Situbondo East Java, Indonesia. *Journal of Pharmacy Research*, 6(7), 685–689. <https://doi.org/10.1016/j.jopr.2013.07.001>
- Agarwal, G. P. (2018). Extraction Techniques for Natural Products. *Journal of Pharmacognosy and Phytochemistry*.
- Andinata, B., Bachtiar, A., Oktamianti, P., Partahi, J. R., & Dini, M. S. A. (2023). A Comparison of Cancer Incidences Between Dharmais Cancer Hospital and GLOBOCAN 2020: A Descriptive Study of Top 10 Cancer Incidences. *Indonesian Journal of Cancer*, 17(2), 119. <https://doi.org/10.33371/ijoc.v17i2.982>
- Aoki, S., Kong, D., Suna, H., Sowa, Y., Sakai, T., Setiawan, A., & Kobayashi, M. (2006). Aaptamine, a spongean alkaloid, activates p21 promoter in a p53-independent manner. *Biochemical and Biophysical Research Communications*, 342(1), 101–106. <https://doi.org/10.1016/j.bbrc.2006.01.119>
- Badar, M. S., Shamsi, S., Ahmed, J., & Alam, Md. A. (2022). *Molecular Dynamics Simulations: Concept, Methods, and Applications*. August, 131–151. https://doi.org/10.1007/978-3-030-94651-7_7
- Baliyan, S., Mukherjee, R., Priyadarshini, A., Vibhuti, A., Gupta, A., Pandey, R. P., & Chang, C. M. (2022). Determination of Antioxidants by DPPH Radical Scavenging Activity and Quantitative Phytochemical Analysis of *Ficus religiosa*. *Molecules*, 27(4). <https://doi.org/10.3390/molecules27041326>
- Bashari, M. H., Agung, M. U. K., Ariyanto, E. F., Al Muqarrabun, L. M. R., Salsabila, S., Chahyadi, A., Rosandy, A. R., Afifah, E., Afni, M., Atmaja, H. E., Putri, T., Utami, F., Subhan, B., Syafrizayanti, Andriani, Y., & Elfahmi. (2025). Two Novel Compounds Isolated from the Marine Fungal Symbiont of *Aspergillus unguis* Induce Apoptosis and Cell Cycle Arrest in Breast Cancer Cells: In vitro Study. *Journal of Experimental Pharmacology*, 17, 37–50. <https://doi.org/10.2147/JEP.S494777>
- Bashari, M. H., Arsydinilhuda, F. Z., Ilhamsyah, R. S., Nugrahani, A. D., Nurdin, R. A., Kartikasari, A., Huda, F., Abdurahman, M., Putri, T., Qomarilla, N., Atmaja, H., Sudji, I. R., Subhan, B., Usman, H. A., Pamela, Y., Ariyanto, E. F., & Meiyanto, E. (2021). The Ethanol Extract of Marine Sponge *Aaptos suberitoides* Suppress Cell Viability, Cell Proliferation and Cell Migration in HER2-Positive Breast Cancer Cell Line. *Asian Pacific Journal of Cancer Prevention*, 22(Supplement 1), 25–32. <https://doi.org/10.31557/APJCP.2021.22.S1.25>
- Bayani, F., Muhali, M., Yuliana, D., Hulyadi, H., & Gargazi, G. (2024). Review of Secondary Metabolites From Melandean Bark Extract (*Bridellia micrantha*): Bioactive Potential and Applications in Health. *Hydrogen: Jurnal Kependidikan Kimia*, 12(3), 413–428. <https://doi.org/10.33394/hjkk.v12i3.11956>

- 22
Bray Bsc, F., Laversanne, I Mathieu, Hyuna, I, Phd, S., Ferlay, J., Siegel Mph, R. L., Soerjomataram, I., Ahmedin, I. & Dvm, J. (2024). *Global cancer statistics 2022: GLOBOC estimates of incidence and mortality worldwide for 36 cancers in 185 countries*. <https://doi.org/10.3322/caac.21834>
- Clark James H., et al. (2017). *Natural Product Extraction Principles and Applications*. <https://doi.org/10.1039/9781849737579-fp001>
- Dyshlovoy, S. A., Fedorov, S. N., Shubina, L. K., Kuzmich, A. S., Bokemeyer, C., Keller-von Amsberg, G., & Honecker, F. (2014). Aaptamines from the marine sponge *Aaptos* sp. display anticancer activities in human cancer cell lines and modulate AP-1-, NF- κ B-, and p53-dependent transcriptional activity in mouse JB6 Cl41 cells. *BioMed Research International*, 2014, 469309. <https://doi.org/10.1155/2014/469309>
- 57
Egbuna, C., Ifemeje, J. C., Udedi, S., & Kumar, S. (2020). *Introduction to Phytochemistry*. CRC Press: Taylor & Francis Group. <https://doi.org/DOI: 10.13140/RG.2.2.18812.2368>
- 1
Fristiohady, A., Sadarun, B., Wahyuni, W., Malaka, M. H., Ahmad, F., Malik, F., Purnama, L. O. M. J., & Sahidin, I. (2020). Isolation and identification of secondary metabolite acetone extract *Aaptos* sp. and its antioxidant properties and acute toxicity. *Journal of Applied Pharmaceutical Science*, 10(6), 81–89. <https://doi.org/10.7324/JAPS.2020.10611>
- 34
Min, H. Y., & Lee, H. Y. (2022). Molecular targeted therapy for anticancer treatment. *Experimental and Molecular Medicine*, 54(10), 1670–1694. <https://doi.org/10.1038/s12276-022-00864-3>
- 30
Molnár, I., Gibson, D., & Krasnoff, S. (2010). Secondary Metabolites from Entomopathogenic Hypocrealean Fungi. *Natural Product Reports*, 27, 1241–1275. <https://doi.org/10.1039/c001459c>
- 23
Nevalainen, H., Kautto, L., & Te'o, J. (2014). Methods for isolation and cultivation of filamentous fungi. *Methods in Molecular Biology (Clifton, N.J.)*, 1096, 3–16. https://doi.org/10.1007/978-1-62703-712-9_1
- 20
Nursid, M. N., Manulang, M., Samiadji, J., & Marraskuranto, E. (2015). Effect of Agitation Speed and Cultivation Time on the Production of the Emestrin Produced by *Emericella nidulans* Marine Fungal. *Squaleen Bulletin of Marine and Fisheries Postharvest and Biotechnology*, 10(2), 73. <https://doi.org/10.15578/squaleen.v10i2.122>
- 27
Oliveira, L. R., Gonçalves, A. R., Quintela, E. D., Colognese, L., Cortes, M. V. de C. B., & de Filippi, M. C. (2024). Genetic Engineering of Filamentous Fungi: Prospects for Obtaining Fourth-Generation Biological Products. In *Applied Microbiology* (Vol. 4, Issue 2, pp. 794–810). <https://doi.org/10.3390/applmicrobiol4020055>
- 16
Oogarah, P. N., Ramanjooloo, A., Doorga, J., Meyepa, C., Van Soest, R., & Marie, D. (2020). Assessing Antioxidant Activity and Phenolic Content of Marine Sponges from Mauritius Waters. *International Journal of Pharmacognosy and Phytochemical Research*, 12(3), 123–131. <https://doi.org/10.25258/phyto.12.3.1>
- 19
Pham, C. D., Hartmann, R., Müller, W. E. G., De Voogd, N., Lai, D., & Proksch, P. (2013). Aaptamine derivatives from the Indonesian sponge *Aaptos suberitoides*. *Journal of Natural Products*, 76(1), 103–106. <https://doi.org/10.1021/np300794b>
- Rendowaty, A., Djamaan, A., & Handayani, D. (2017). Waktu Kultivasi Optimal dan Aktivitas Antibakteri dari Ekstrak Etil Asetat Jamur Simbion *Aspergillus unguis* (WR8) dengan *Haliclona fascigera*. *Jurnal Sains Farmasi & Klinis*, 4(1), 49. <https://doi.org/10.29208/jsfk.2017.4.1.147>

- Rivera, A. P., & Uy, M. M. (2012). In vitro antioxidant and cytotoxic activities of some marine sponges collected off misamis oriental coast, Philippines. *E-Journal of Chemistry*, 9(1), 354–358. <https://doi.org/10.1155/2012/506046>
- Sajna, K. V., Kamat, S., & Jayabaskaran, C. (2020). Antiproliferative Role of Secondary Metabolites From *Aspergillus unguis* AG 1.1 (G) Isolated From Marine Macroalgae *Enteromorpha* sp. by Inducing Intracellular ROS Production and Mitochondrial Membrane Potential Loss Leading to Apoptosis. *Frontiers in Marine Science*, 7. <https://doi.org/10.3389/fmars.2020.543523>
- Santaniello, G., Nebbioso, A., Altucci, L., & Conte, M. (2023). Recent Advancement in Anticancer Compounds from Marine Organisms: Approval, Use and Bioinformatic Approaches to Predict New Targets. *Marine Drugs*, 21(1). <https://doi.org/10.3390/md21010024>
- Situmorang, Fronika, N. (2012). Uji Toksisitas Ekstrak Etanol Spons *Aaptos suberitoides* Menggunakan Larva Udang *Artemia Salina* Leach. *Fakultas Farmasi Universitas Pancasila*.
- Stange, S., Steudler, S., Delenk, H., Werner, A., Walther, T., & Wagenführ, A. (2019). Influence of Environmental Growth Factors on the Biomass and Pigment Production of *Halorociboria aeruginascens*. *Journal of Fungi (Basel, Switzerland)*, 5(2). <https://doi.org/10.3390/jof5020046>
- Tailor, N. K., Deswal, G., & Grewal, A. S. (2024). Aaptamine: A Versatile Marine Alkaloid for antioxidant, Antibacterial, and Anticancer Therapeutics. *Chemistry (Switzerland)*, 6(4), 677–694. <https://doi.org/10.3390/chemistry6040040>
- Turon, M., J., U. M., & Daniel, M. (2019). Multipartner Symbiosis across Biological Domains: Looking at the Eukaryotic Associations from a Microbial Perspective. *MSystems*, 4(4), 10.1128/msystems.00148-19. <https://doi.org/10.1128/msystems.00148-19>
- Van Anh, C., Kang, J. S., Choi, B. K., Lee, H. S., Heo, C. S., & Shin, H. J. (2021). Polyketides and meroterpenes from the marine-derived fungi *aspergillus unguis* 158SC-067 and *A. Flocculosus* 01NT-1.1.5 and their cytotoxic and antioxidant activities. *Marine Drugs*, 19(8). <https://doi.org/10.3390/md19080415>
- Varijakzhan, D., Loh, J.-Y., Yap, W.-S., Yusoff, K., Seboussi, R., Lim, S.-H. E., Lai, K.-S., & Chong, C.-M. (2021). Bioactive Compounds from Marine Sponges: Fundamentals and Applications. *Marine Drugs*, 19(5). <https://doi.org/10.3390/md19050246>
- Yun, C. W., Kim, H. J., & Lee, S. H. (2019). Therapeutic Application of Diverse Marine-derived Natural Products in Cancer Therapy. *Anticancer Research*, 39(10), 5261–5284. <https://doi.org/10.21873/anticancer.13721>
- Zakwan, M., Sutriana, A., Nurliana, Asmiliana, N., Ammar, M., & Novianti, A. N. (2023). Toxicity Test of Flavonoid Compounds from Ethyl Acetate Extract of Malacca Leaves with Brine Shrimp Lethality Test. *Jurnal Medik Veteriner*, 6(3), 326–330. <https://doi.org/10.20473/jmv.vol6.iss3.2023.326-330>

ORIGINALITY REPORT

42%

SIMILARITY INDEX

37%

INTERNET SOURCES

36%

PUBLICATIONS

26%

STUDENT PAPERS

PRIMARY SOURCES

1

ouci.dntb.gov.ua

Internet Source

2%

2

repository.usd.ac.id

Internet Source

2%

3

www.mdpi.com

Internet Source

2%

4

impactfactor.org

Internet Source

1%

5

www.bioflux.com.ro

Internet Source

1%

6

Vinita Manimaran Nadar, Selvambigai Manivannan, Rajasekar Chinnaiyan, Muthusamy Govarthanan, Kumar Ponnuchamy. "Review on marine sponge alkaloid, aaptamine: A potential antibacterial and anticancer drug", Chemical Biology & Drug Design, 2021

Publication

1%

7

Submitted to University of Keele

Student Paper

1%

8

ejournal.sisfokomtek.org

Internet Source

1%

9

repository.ub.ac.id

Internet Source

1%

10

e-journal.unair.ac.id

Internet Source

1%

| | | |
|----|---|-----|
| 11 | Muhammad Zakwan, Amalia Sutriana, Nurliana Nurliana, Nuzul Asmilia, Muhammad Ammar, Arindita Niatazya Novianti. "Toxicity Test of Flavonoid Compounds from Ethyl Acetate Extract of Malacca Leaves with Brine Shrimp Lethality Test", Jurnal Medik Veteriner, 2023 Publication | 1 % |
| 12 | Submitted to Sefako Makgatho Health Science University Student Paper | 1 % |
| 13 | Trini Suryowati, John Jackson Yang, Lusia Sri Sunarti, Maria Bintang. "The Effect of Torbangun (Coleus amboinicus Lour) Leaf Extract on Antibiotic Resistant Bacteria Escherichia coli pBR322 and Toxicity Tests on Artemia salina Leach", Indonesian Journal of Applied Research (IJAR), 2024 Publication | 1 % |
| 14 | Submitted to University of Portsmouth Student Paper | 1 % |
| 15 | journal-center.litpam.com Internet Source | 1 % |
| 16 | ph04.tci-thaijo.org Internet Source | 1 % |
| 17 | www.indonesianjournalofcancer.or.id Internet Source | 1 % |
| 18 | pubmed.ncbi.nlm.nih.gov Internet Source | 1 % |
| 19 | assets-eu.researchsquare.com Internet Source | 1 % |
| 20 | Submitted to The University of the West of Scotland | 1 % |

| | | |
|----|--|-----|
| 21 | www.um.edu.mt Internet Source | 1 % |
| 22 | Ana Carolina M. Domingues, Soraia B. de Oliveira, Nayara G. Tessarollo, Ana Paula Lepique et al. "Use of patient-derived organotypic tumor spheroids for testing of viral vector gene therapy in combination with checkpoint blockade", Molecular Therapy Oncology, 2025 Publication | 1 % |
| 23 | media.suub.uni-bremen.de Internet Source | 1 % |
| 24 | Submitted to Dicle University Student Paper | 1 % |
| 25 | Submitted to University of the Philippines - Main Library Student Paper | 1 % |
| 26 | Submitted to Queen Mary and Westfield College Student Paper | 1 % |
| 27 | www.preprints.org Internet Source | 1 % |
| 28 | doaj.org Internet Source | 1 % |
| 29 | Nanda Putra, Afrillia Nuryanti Garmana, Nurinanda Prisky Qomaladewi, Amrianto et al. "Bioactivity-guided isolation of a bioactive compound with α -glucosidase inhibitory activity from the leaves extract of Sauropus androgynus", Sustainable Chemistry and Pharmacy, 2023 Publication | 1 % |

| | | |
|----|--|------|
| 30 | www.researchsquare.com Internet Source | 1 % |
| 31 | "Natural Secondary Metabolites", Springer Science and Business Media LLC, 2023 Publication | 1 % |
| 32 | Submitted to Flinders University Student Paper | 1 % |
| 33 | journals.sbm.u.ac.ir Internet Source | 1 % |
| 34 | Submitted to Uludag University Student Paper | 1 % |
| 35 | Submitted to Taylor's Education Group Student Paper | 1 % |
| 36 | Iis Yusma Fitri, Kurniawanti Kurniawanti, Gustini Syahbirin, Purwantiningsih Sugita. "Antioxidant Activity, Cytotoxicity, and Identification of Secondary Metabolites of <i>Kigelia africana</i> from Waterpark Platinum Riau", Indo. J. Chem. Res., 2023 Publication | <1 % |
| 37 | Zahra Izadiyan, Misni Misran, Katayoon Kalantari, Thomas Webster, Pooneh Kia, Noor Basrowi, Elisa Rasouli, Kamyar Shameli. "Advancements in Liposomal Nanomedicines: Innovative Formulations, Therapeutic Applications, and Future Directions in Precision Medicine", International Journal of Nanomedicine, 2025 Publication | <1 % |
| 38 | peerj.com Internet Source | <1 % |
| 39 | Submitted to Universitas Tanjungpura Student Paper | <1 % |

| | | |
|----|--|------|
| 40 | mdpi-res.com Internet Source | <1 % |
| 41 | www.dovepress.com Internet Source | <1 % |
| 42 | bestjournal.untad.ac.id Internet Source | <1 % |
| 43 | www.sci.ku.ac.th Internet Source | <1 % |
| 44 | zenodo.org Internet Source | <1 % |
| 45 | Muhammad Hasan Bashari, Mochamad Untung Kurnia Agung, Eko Fuji Ariyanto, Laode Muhammad Ramadhan Al Muqarrabun et al. "Two Novel Compounds Isolated from the Marine Fungal Symbiont of <i>Aspergillus unguis</i> Induce Apoptosis and Cell Cycle Arrest in Breast Cancer Cells: In vitro Study", <i>Journal of Experimental Pharmacology</i> , 2025 Publication | <1 % |
| 46 | Submitted to Universitas Diponegoro Student Paper | <1 % |
| 47 | inscricao.faculdadeitop.edu.br Internet Source | <1 % |
| 48 | phcogj.com Internet Source | <1 % |
| 49 | Basma Hossam Abdelmonem, Lereen T. Kamal, Rana M. Elbaz, Mohamed R. Khalifa, Anwar Abdelnaser. "From contamination to detection: The growing threat of heavy metals", <i>Heliyon</i> , 2025 Publication | <1 % |

| | | |
|----|---|------|
| 50 | Submitted to University of Northumbria at Newcastle Student Paper | <1 % |
| 51 | Submitted to Universiti Malaysia Sabah Student Paper | <1 % |
| 52 | www.growkudos.com Internet Source | <1 % |
| 53 | "Antibacterial Activity of Indonesian Sponge Associated Fungi Against Clinical Pathogenic Multidrug Resistant Bacteria", Journal of Applied Pharmaceutical Science, 2018 Publication | <1 % |
| 54 | Submitted to Queen's University of Belfast Student Paper | <1 % |
| 55 | perpus.univpancasila.ac.id Internet Source | <1 % |
| 56 | Submitted to Mt. San Jacinto College Student Paper | <1 % |
| 57 | etd.repository.ugm.ac.id Internet Source | <1 % |
| 58 | journal.unimma.ac.id Internet Source | <1 % |
| 59 | www.scientific.net Internet Source | <1 % |
| 60 | Submitted to Unviersidad de Granada Student Paper | <1 % |
| 61 | Zahro Rizka Shafira, Susi Endrini, Wan Nor I'zzah Wan Mohamad Zain, Nur Syafina Binti Hisham. "Cytotoxic Effects and Anticancer Activity Assay of Sonneratia alba Fruit on Cancer Cell Cultures CaCo-2 and MCF-7", Asian Journal of Biological Sciences, 2024 | <1 % |

| | | |
|----|---|------|
| 62 | mafiadoc.com Internet Source | <1 % |
| 63 | tjnpr.org Internet Source | <1 % |
| 64 | Submitted to Griffith University Student Paper | <1 % |
| 65 | Lourdes Inés Pascual, Lorena Luna, Roxana Elizabeth González, Javier Esteban Ortiz et al. "Correlation Analysis Among the Chemical Composition and Cytotoxic and Antioxidative Activities of a Tessaria absinthioides Decoction for Endorsing Its Potential Application in Oncology", Plants, 2024 Publication | <1 % |
| 66 | dom.lndb.lv Internet Source | <1 % |
| 67 | issuu.com Internet Source | <1 % |
| 68 | link.springer.com Internet Source | <1 % |
| 69 | science.rsu.lv Internet Source | <1 % |
| 70 | simakip.uhamka.ac.id Internet Source | <1 % |
| 71 | Afsona Khatun, Kuntal Hazra, Belarani Mahato, Ritabrata Koley et al. "Studies of chemical distribution and pharmacological activities of porifera-derived alkaloids: A review (2000–2023)", European Journal of Medicinal Chemistry Reports, 2024 Publication | <1 % |

| | | |
|----|--|------|
| 72 | Idin Sahidin, Wahyuni, Arief Rabbani Rahim, Muhammad Arba et al. "Radical scavenging and antimicrobial activities of diarylheptanoids and steroids from Etlingera calophrys rhizome", Sustainable Chemistry and Pharmacy, 2022 Publication | <1 % |
| 73 | Musdalipah Musdalipah, Selfyana Austin Tee, Karmilah Karmilah, Sahidin Sahidin, Adryan Fristiohady, Agung Wibawa Mahatva Yodha. "Total Phenolic and Flavonoid Content, Antioxidant, and Toxicity Test with BSLT of Meistera chinensis Fruit Fraction from Southeast Sulawesi", Borneo Journal of Pharmacy, 2021 Publication | <1 % |
| 74 | www.ncbi.nlm.nih.gov Internet Source | <1 % |
| 75 | www.researchgate.net Internet Source | <1 % |
| 76 | "Microbial Biocontrol: Food Security and Post Harvest Management", Springer Science and Business Media LLC, 2022 Publication | <1 % |
| 77 | "Nano-formulation of Dietary Phytochemicals for Cancer Management", Springer Science and Business Media LLC, 2024 Publication | <1 % |
| 78 | A. P. Rivera, M. M. Uy. " Antioxidant and Cytotoxic Activities of Some Marine Sponges Collected off Misamis Oriental Coast, Philippines ", E-Journal of Chemistry, 2012 Publication | <1 % |

- | | | |
|----|---|------|
| 79 | Elin Julianti, Muhammad Azhari, Marlia Singgih, Fitri Siti Dzurriyyah Aminah et al. "Cytotoxic activity of cordycepin produced by marine-derived fungus Emericella sp. against HT29 human colon cancer cell lines", Archives of Microbiology, 2023 Publication | <1 % |
| 80 | Jun-Ping Shiau, Min-Yu Lee, Jen-Yang Tang, Hsin Huang et al. "Marine Sponge Aaptos suberitoides Extract Improves Antiproliferation and Apoptosis of Breast Cancer Cells without Cytotoxicity to Normal Cells In Vitro", Pharmaceuticals, 2022 Publication | <1 % |
| 81 | Michael Qwarse, Alphonse Ignace Marealle, Francis Machumi, Matobola Joel Mihale et al. "Exploring the Therapeutic Potential of Wild Edible Mushrooms: Safety Evaluation and Isolation of Antimycobacterial Sterols from Afrocantharellus platyphyllus (Heinem.) Tibuhwa", Chemistry Africa, 2023 Publication | <1 % |
| 82 | idoc.pub Internet Source | <1 % |
| 83 | iieta.org Internet Source | <1 % |
| 84 | "Fungal Biopolymers and Biocomposites", Springer Science and Business Media LLC, 2022 Publication | <1 % |
| 85 | Rahmi Vika Ulia, Suryati Suryati, Adlis Santoni. "Cytotoxic Potential of Essential Oil Isolated from Clibadium Surinamese L Leaves Against | <1 % |

T47D Breast and HeLa Cervical Cancer Cells",
Molekul, 2023

Publication

86

Hb25_Springer Handbook of Marine
Biotechnology, 2015.

Publication

<1 %

87

Shinta R. Dewi, Lee A. Stevens, Muslih Anwar,
Yujie Mao, Rebecca S. Ferrari, Derek J. Irvine,
Eleanor R. Binner. "Towards a scalable cacao
pod husk biorefinery: Understanding the
effects of process conditions on phenolic
antioxidant extraction and residual solid
properties", Chemical Engineering Science,
2025

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off